



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/54, 1/26, 1/28	A1	(11) International Publication Number: WO 92/22669 (43) International Publication Date: 23 December 1992 (23.12.92)
<p>(21) International Application Number: PCT/GB92/01116</p> <p>(22) International Filing Date: 19 June 1992 (19.06.92)</p> <p>(30) Priority data: 9113212.6 19 June 1991 (19.06.91) GB</p> <p>(71) Applicant (for all designated States except US): HYPO-GUARD (UK) LIMITED [GB/GB]; Dock Lane, Melton, Woodbridge, Suffolk IP12 1PE (GB).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only) : GULLICK, Stephen, Peter [GB/GB]; 18 Deben Avenue, Martlesham, Ipswich, Suffolk IP5 7PQ (GB).</p> <p>(74) Agent: DUMMETT, Thomas, Ian, Peter; Dummett Copp & Co., 14 The Square, Martlesham Heath, Ipswich, Suffolk IP5 7SL (GB).</p>	<p>(81) Designated States: AT, AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE, DE (European patent), DK, DK (European patent), ES, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL, NL (European patent), NO, PL, RO, RU, SD, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: REAGENT MIXTURES FOR GLUCOSE ASSAY</p> <p>(57) Abstract</p> <p>The present invention relates to a test reagent mixture composition comprising the enzymes glucose oxidase and peroxidase and a chromogen which interacts with the hydrogen peroxide from the oxidation of the blood glucose by the glucose oxidase, characterised in that the glucose oxidase and the peroxidase are present in proportions which provide from 300 to 700 International Units (IUs) of glucose oxidase and at least 20 International Units of peroxidase and in that the chromogen is present in an amount which provides from 12 to 20 grams of active chromogen per 500,000 International Units of glucose oxidase. Preferably, the composition is put up in a low molecular weight gelatin matrix and is impregnated into a micro-porous carrier membrane.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LJ	Lichtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MC	Malagascar		
ES	Spain				

REAGENT MIXTURES FOR GLUCOSE ASSAY

The present invention relates to a reagent mixture, notably to a mixture of analytical reagents in a carrier gel which provides enhanced consistency of the colour generated with the elapse of time.

BACKGROUND TO THE INVENTION:

10 Blood samples are often assessed for the amount of glucose or some other constituent therein by reacting the blood with one or more reagents carried on a test stick or pad so as to develop a colour which can be observed by an operator. For example, a reagent pad can contain the enzymes glucose oxidase and peroxidase and o-tolidine as the chromogen which turns blue as the glucose oxidase oxidises glucose in the blood sample to gluconic acid and hydrogen peroxide. The hydrogen peroxide reacts in the presence of the peroxidase with the o-tolidine to give a blue colour whose intensity depends upon the amount of hydrogen peroxide released and hence the amount of glucose in the blood. The reagents are usually put up in a gel matrix, for example of a natural gel, for example a gelatin, or of a synthetic polymer, for example a polyvinylpyrrolidone.

25 However, problems arise in that the colour is affected by the time over which the blood sample is held in contact with the reagent pad, as well as the amount of blood in contact with the reagents. It is therefore customary for such tests to be carried out within a strictly monitored time schedule and the results are often of dubious value due to inaccuracies in observing the time schedule.

35 Surprisingly, we have found that the proportion of the reagents to one another in the matrix affects the period over which a consistent colour is produced by the interaction of the blood with the reagent. If the proportions in the mixture lie within

certain limits, the colour produced is sufficiently constant over a period of time for the need for strict adherence to a time schedule to be reduced.

5 SUMMARY OF THE INVENTION:

Accordingly, the present invention provides a blood test reagent mixture composition comprising the enzymes glucose oxidase and peroxidase and a chromogen which interacts with the
10 hydrogen peroxide from the oxidation of the blood glucose by the glucose oxidase, characterised in that the glucose oxidase and the peroxidase are present in proportions which provide from 300 to 700 International Units (IUs) of glucose oxidase and at least 20 International Units of peroxidase and in that
15 the chromogen is present in an amount which provides from 12 to 20 grams of active chromogen per 500,000 International Units of glucose oxidase. Preferably, the glucose oxidase is present in from 400 to 550 IUs per 27.5 to 32.5 IUs of peroxidase and the
20 chromogen is o-tolidine which is present in an amount of from 12 to 15 gs per 500,000 IUs of the glucose oxidase.

It is preferred that the reagent mixture be put up in a gel matrix, notably a gelatin matrix, which provides from 200 to 400 gs of the matrix on a dry weight basis per 500,000 IUs of
25 the glucose oxidase.

It is particularly preferred that the reagent mixture/matrix be absorbed in a micro-porous membrane carrier.

30 The enzymatic reagents as used herein can be present in any suitable form, for example as the dry powdered active enzyme or as a precursor or addition product thereof which under the conditions of the test to be carried out produces an active enzyme in the reagent mixture. Thus, the enzymatic reagent can
35 be an active enzyme, for example glucose oxidase or peroxidase, or a stabilised form thereof, for example an acetate or other

salt or adduct thereof, which releases the active enzyme when the reagent mixture is wetted.

For convenience, the invention will be described hereinafter in terms of a mixture of glucose oxidase and peroxidase as conventionally used in the assessment of glucose in a blood sample.

Similarly, the term chromogenic material is used herein to denote any material which develops a property upon interaction with one or more of the products produced when the enzymatic reagent reacts with the blood sample to be assessed. Thus, the term includes materials which develop ultraviolet fluorescence or other detectable but not visible properties. However, it is preferred that the chromogenic material be one which develop a colour within the visible spectrum, for example as when di-anisidine or o-tolidine reacts with the hydrogen peroxide released when glucose in blood interacts with the glucose oxidase in the reagent mixture. The chromogenic material can be used in the form of the active material or a precursor or adduct thereof, notably an inorganic acid salt thereof such as the hydrochloride or sulphate, which releases the active ingredient during the test.

For convenience, the invention will be described hereinafter in terms of o-tolidine as the chromogenic material.

The enzymatic reagent and chromogenic material are operatively associated with one another so that they can interact under the conditions of the test procedure. Typically, the reagent and the chromogenic material will be put up in physical admixture with one another. However, it is within the scope of the present invention to put up the reagent and chromogenic material in a two part form which is admixed immediately prior to use; or in a form in which the reaction products of the interaction of the enzyme reagent with the glucose in the blood

sample diffuse into a zone containing the chromogenic material to develop the colour therewith separately from the enzyme interaction zone. Thus, for example, the enzyme reagent can be concentrated at one end or one side of a reagent pad and the
5 chromogenic material at the other end or side.

For convenience, the invention will be described hereinafter in terms of a pad of the reagent mixture containing the enzymatic reagent and the chromogenic material substantially uniformly
10 distributed throughout the pad.

The reagent mixture may contain other materials as is customary, for example phosphate buffering agents, preservatives, anti-coagulants or surface active agents. Such
15 other materials are typically inert to the material to be tested, the other constituents of the reagent mixture and the products of the reactions and interactions which occur during the test procedure. Such other constituents can be present in the amounts normally used in such reagent mixtures.

20

As stated above, we have found that if the enzymatic reagent and chromogenic material are present in the reagent mixture composition within specified proportions, the colour which the interactions between the material being assessed and the
25 various components of the mixture is surprisingly stable and enables the colour to be observed over a wider period of time than hitherto. Thus, the enzymatic reagents will typically be present in proportions of from 400 to 600, notably 450 to 550, International Units (IUs) of glucose oxidase and at least 20
30 IUs, typically about 27.5 to 35 IUs of peroxidase in the mixture. The chromogenic material will typically be present in an amount of from 12 to 17, notably from 13 to 16, grams per 500,000 IUs of the glucose oxidase. The optimum proportions within these ranges can be determined for any given case and a
35 given carrier by simple trial and error tests.

As stated above, the reagent mixture is preferably put up in a matrix carrier medium so that the material to be assessed can penetrate to the enzyme reagent and the chromogenic material. The matrix can be provided by a natural gum, jelly or gel, for example a gelatin, agar agar, aspic or silica gel; or can be provided by a synthetic polymer gel, for example a cellulosic gel or a polyvinyllic resin gel. For convenience, the invention will be described hereinafter in terms of the use of a gelatin gel as the carrier matrix.

The gel matrix can carry the enzymatic reagent and chromogenic material substantially uniformly distributed throughout it. This is conveniently achieved by mixing the enzyme reagents into a premix of the gelling agent and the chromogenic material; and allowing the mixture to set in the desired form. Alternatively, the enzyme reagent and the chromogenic material can be admixed with a thixotropic gel carrier which is worked, for example by being stirred, to maintain it in the fluid state during mixing, but which is then allowed to set for storage and transport prior to use.

Alternatively, the matrix may contain the enzymatic reagent and chromogenic material non-uniformly distributed therein, as when a gel layer is formed which has a high concentration of the gel matrix in its upper layers to act a protective layer or coating for the reagent rich lower layers; or where the enzymatic reagent is located in a separate zone of the matrix from that containing the chromogenic material. In this case, the product from the interaction of the material being tested with one or more of the enzymatic reagents diffuses from the enzyme zone into the zone containing the chromogenic material to develop a colour as a separate stage in that zone.

For convenience, the invention will be described hereinafter in terms of a reagent mixture which is uniformly distributed throughout a gelatin matrix.

The matrix carrying the reagent mixture can be put up in a number of physical forms, for example as test strips or discs in which a pad of the matrix is applied to one face of the strip or disc and the colour resulting from the interaction of the material under assessment and the reagents and materials in the matrix is observed visually against the white background of the support strip or disc or against a separate reference background. Alternatively, the matrix can be put up in a series of zones through which the reagent mixture is distributed so that the interaction of the material to be assessed with the enzyme occurs in one zone and the product of that interaction diffuses to a second zone in which the colour reaction takes place. In a further alternative, the reagent mixture matrix can be absorbed or impregnated into the pores of a porous carrier to one face of which the material to be assessed is applied and the colour developing within the matrix is viewed from the opposite surface of the carrier.

For convenience, the invention will be described in terms of the use of a pad or disc of a micro-porous membrane which is impregnated with the reagent matrix.

In a conventional blood test reagent mixture, the gel matrix is a high molecular weight gelatin which is present in about 4% by dry weight. However, where the reagent mixture is to be absorbed into a micro-porous membrane, we prefer to use a low molecular weight gelatin, typically with a molecular weight in the range 20,000 to 50,000. Where such a gelatin is present in the amounts used hitherto, we have found that this results in a gel matrix which cannot be held satisfactorily within the pores of the membrane. On the other hand, we have found that if the gel content of the reagent mixture composition exceeds about 20% by dry weight, the gel inhibits the diffusion of reaction products through the membrane and hence development of a colour reflecting the true extent of the interactions which have occurred. We therefore prefer to provide the gel matrix

as a low molecular weight gelatin in an amount of from 250 to 325 gs by dry weight per 500,000 IUs of the glucose oxidase present in the reagent mixtrue.

- 5 The invention will now be illustrated by the following Example in which all parts and percentages are given by weight unless stated otherwise.

10 A first solution was made by stirring together at room temperature 300 mls of de-ionised water, 200 mls of 0.5 Molar sodium phosphate buffer to give a pH of 7, 100 mls of a 20% w/v solution of the surfactant Gantrez and 300 gs of dry powdered gelatin having a molecular weight in the range 25,000 to 40,000.

15 A second solution was prepared by stirring together at 60° C for one hour 300 mls of de-ionised water, 300 mls of methoxyethanol and 15 gs of o-tolidine hydrochloride or dianisidine hydrochloride.

20 The second solution was mixed dropwise with stirring into the first solution and the mixture stood for 1 hour at 60° C.

25 A third solution was made up by mixing 500,000 IUs of glucose oxidase and 30,000 IUs of peroxidase in a 0.1 Molar solution of the sodium phosphate buffer. This solution was mixed with stirring into the other mixed solutions and filtered through a 0.1 micrometre aperture filter.

30 The resultant solution was impregnated into a polysulfone resin sheet (0.2 to 0.4 mms thick and having an average pore diameter of 0.2 micrometres and an air permeability of 3 litres per minute per square centimetre at an applied pressure of 10 psig) to provide 5 IUs of glucose oxidase, 3 IUs of peroxidase, 0.2
35 milligrams of o-tolidine and 4 milligrams of gelatin per square centimetre of the membrane.

By way of comparison, the same membrane was impregnated with a conventional reagent mixture to provide the conventional level of enzyme and chromogen per square centimetre.

5 Blood samples were applied to the faces of a number of 6 mms diameter discs cut from each of the membranes. With the reagent compositions of the invention, a blue colour developed after only 10 seconds. The hue and intensity of the colour became stable after about 30 to 40 seconds and remained stable
10 for a further 30 to 40 seconds, thus allowing considerable latitude for the time to observe the colour. By way of comparison, the conventional formulations gave a colour which deepened in hue and intensity over 10 to 30 seconds after applying the blood sample, but which degenerated after a
15 further 15 seconds, giving little or no latitude in the time for observing the true colour.

From another aspect, the present invention provides a method for making a test reagent mixture of the invention, wherein the
20 components of the mixture are admixed with one another to provide a substantially uniform mixture of the components.

The invention further provides a method for making a test reagent mixture carried on a micro-porous carrier medium,
25 wherein a fluid reagent mixture of the invention is applied to the carrier medium. Preferably, the mixture is applied by impregnating the medium, for example by padding a sheet of the carrier through a bath of the reagent mixture, and allowing the mixture to gel within the pores of the carrier. Preferably,
30 the gelled mixture blinds the bores of the pores of the carrier so that rupture of blood or other cells due to capillary action by the pores is reduced. Discs or other shapes can be cut from the impregnated carrier for mounting on tests sticks having apertures therein or as the end walls of sample receivers so
35 that the colour which develops in the carrier can be observed from the opposite side to that to which the blood is applied.

CLAIMS:

1. A blood test reagent mixture composition comprising the enzymes glucose oxidase and peroxidase and a chromogen which
5 interacts with the hydrogen peroxide from the oxidation of the blood glucose by the glucose oxidase, characterised in that the glucose oxidase and the peroxidase are present in proportions which provide from 300 to 700 International Units (IUs) of
10 glucose oxidase and at least 20 International Units of peroxidase and in that the chromogen is present in an amount which provides from 12 to 20 grams of active chromogen per 500,000 International Units of glucose oxidase.
2. A test reagent mixture as claimed in claim 1,
15 characterised in that the glucose oxidase is present in from 400 to 550 IUs per 27.5 to 32.5 IUs of peroxidase and the chromogen is o-tolidine which is present in an amount of from 12 to 17 gs per 500,000 IUs of the glucose oxidase.
- 20 3. A test reagent mixture as claimed in either of claims 1 or 2, characterised in that it is put up in a gel matrix.
4. A test reagent mixture as claimed in claim 3,
25 characterised in that the gel matrix is a gelatin matrix, which provides from 200 to 400 gs of gelatin on a dry weight basis per 500,000 IUs of the glucose oxidase.
5. A test reagent mixture as claimed in any one of the preceding claims, characterised in that the reagent mixture is
30 carried by a micro-porous membrane.
6. A test reagent mixture as claimed in either of claims 4 or 5, characterised in that the gelatin has a molecular weight in the range 20,000 to 50,000 and is present in an amount of from
35 250 to 325 gs by dry weight per 500,000 IUs of the glucose oxidase.

7. A test reagent mixture according to claim 1, substantially as hereinbefore described in the Example.

8. A method for making a test reagent mixture as claimed in claim 1, characterised in that the components of the mixture are admixed with one another to provide a substantially uniform mixture of the components.

9. A method for making a test reagent mixture as claimed in claim 5, characterised in that a fluid reagent mixture as claimed in any one of claims 1 to 4 or claim 6 is impregnated into the pores of a micro-porous carrier membrane.

10. A method for testing blood samples, characterised in that it comprises applying blood to a test reagent mixture as claimed in claim 1 and observing the colour which develops.

11. A method as claimed in claim 10, characterised in that the reagent mixture is carried by a micro-porous carrier membrane and the blood is applied to one face of the membrane and the colour is observed from the opposite face of the membrane.

25

30

35

INTERNATIONAL SEARCH REPORT

PCT/GB 92/01116

International Application No

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12Q1/54; C12Q1/26; C12Q1/28		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12Q	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ⁹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	US,A,4 340 669 (R. BAUER) 20 July 1982 see the whole document	1-11
A	EP,A,0 016 947 (BOEHRINGER MANNHEIM GMBH.) 15 October 1980 see the whole document	1-11
A	FR,A,2 295 425 (BOEHRINGER MANNHEIM GMBH.) 16 July 1976 see the whole document	1-11
A	GB,A,893 318 (MILES LABORATORIES, INC.) 4 April 1962 see the whole document	1-11
<p>⁹ Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
02 OCTOBER 1992	11 5. 10. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	GRIFFITH G. <i>Gerard Griffith</i>	

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO. GB 9201116
SA 60990**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 02/10/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4340669	20-07-82	AT-T- 9821	15-10-84
		AU-B- 524238	09-09-82
		CA-A- 1157749	29-11-83
		EP-A, B 0058334	25-08-82
		JP-C- 1311966	11-04-86
		JP-A- 57146599	10-09-82
		JP-B- 60035118	13-08-85
		US-A- 4391905	05-07-83
		US-A- 4391906	05-07-83
EP-A-0016947	15-10-80	DE-B- 2913553	09-10-80
		AT-T- 4126	15-07-83
		AU-B- 515851	07-05-81
		AU-A- 5677980	09-10-80
		CA-A- 1152871	30-08-83
		JP-C- 1111586	31-08-82
		JP-A- 55135599	22-10-80
		JP-B- 56046800	05-11-81
		US-A- 4517287	14-05-85
FR-A-2295425	16-07-76	DE-A- 2460903	24-06-76
		AT-B- 350189	10-05-79
		AU-B- 504794	01-11-79
		AU-A- 8755775	23-06-77
		BE-A- 836789	18-06-76
		CA-A- 1060906	21-08-79
		CH-A- 619048	29-08-80
		DE-C- 2462952	01-10-87
		GB-A- 1464360	09-02-77
		GB-A- 1464359	09-02-77
		JP-A- 54036237	16-03-79
		JP-C- 1263517	16-05-85
		JP-A- 51089491	05-08-76
		JP-B- 54003394	22-02-79
		NL-A- 7514628	23-06-76
		SE-B- 458687	24-04-89
SE-A- 7514196	22-06-76		
US-A- 4385114	24-05-83		
GB-A-893318		US-A- 3012976	

EPO FORM P0077

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82